Astronomy & Astrophysics manuscript no.
(DOI: will be inserted by hand later)

February 1, 2008

arXiv:astro-ph/0110426v3 21 Feb 2002

Abundance and evolution of galaxy clusters in cosmological models with massive neutrino

N.A. Arhipova[1], T. Kahniashvili[2,3], and V.N. Lukash[1]

[1] Astro Space Center of P.N.Lebedev Physical Institute, Moscow, Russia
[2] Center for Plasma Astrophysics, Abastumani Astrophysical Observatory, Tbilisi, Georgia
[3] Department of Physics and Astronomy, Rutgers University, Piscataway, NJ, USA

Received 2 August 2001 / accepted 11 January 2002

Abstract. The time evolution of the number density of galaxy clusters and their mass and temperature functions are used to constrain cosmological parameters in the spatially flat dark matter models containing hot particles (massive neutrino) as well as cold and baryonic matter. We test the modified MDM ($\Lambda = 0$) models with cosmic gravitational waves and show that they neither pass the cluster evolution test nor reproduce the observed height of the first acoustic peak in $\Delta T/T$ spectrum, and therefore should be ruled out. The models with a non-zero cosmological constant are in better agreement with observations. We estimate the free cosmological parameters in ΛMDM with a negligible abundance of gravitational waves, and find that within the parameter ranges $h \in (0.6, 0.7)$, $n \in (0.9, 1.1)$, $f_\nu \equiv \Omega_\nu/\Omega_m \in (0, 0.2)$, (i) the value of Ω_Λ is strongly affected by a small fraction of hot dark matter: $0.45 < \Omega_\Lambda < 0.7$ (1σ CL), and (ii) the redshift evolution of galaxy clusters alone reveals the following explicit relation between Ω_Λ and f_ν:

$$\Omega_\Lambda + 0.5 f_\nu = 0.65 \pm 0.1.$$

This degeneracy is also expected in LSS tests (with a smaller error). The present accuracy of observational data allows to bound the fraction of hot matter, $f_\nu \in (0, 0.2)$; the number of massive neutrino species remains undelimited, $N_\nu = 1, 2, 3$.

Key words. cosmology: observations – cosmology: theory – large–scale structure of the Universe – clusters of galaxies

1. Introduction

One of the most important problems of modern cosmology is the formation of large-scale structure in the Universe (LSS). The last decade progress in observations of LSS and cosmic microwave background anisotropy (CMBA) has allowed a productive comparison of theory with observations. Any realistic cosmological model should be consistent with LSS observational data in the range of scales from sub-Mpc (Gnedin 1998, Peackock 1997) up to the cosmological horizon (Smoot et al. 1992, Jaffe et al. 2001).

According to the standard theory of gravitational instability the observable LSS in the Universe has been formed by growth of small density inhomogeneities generated during the very early stages of cosmological expansion. Two main features of the primordial perturbation field – the Gaussian character of linear density perturbations and the bottom-up LSS formation – are confirmed by available astrophysical observations. Therefore, assuming that the initial conditions are given one can derive the theoretical predictions for LSS formation in a dark matter (DM) model and then use statistical analysis and observational data to test the viability of the considered models, i.e. to constrain the allowed range of principal cosmological parameters.

Nowadays, the most popular cosmological models are cold DM with non-zero cosmological constant (ΛCDM) (Peebles 1984, Kofman & Starobinsky 1985) and mixed DM without (MDM) (Fang et al. 1984, Shafi & Stecker 1984, Valdarnini & Bonometto 1984, Lukash 1991, Lucchin et al. 1996) and with (ΛMDM) (Kahniashvili et al. 1996, Novosyadlyj et al. 1999b and refs. therein) cosmological constant, where MDM is in the form of non-baryonic cold (neutralino or hypothetical axions) and hot (massive neutrino) particles.

The cosmological impact of non-baryonic collisionless matter depends on the free-streaming path of DM particles: at small scales ($k > k_{FS}$) the perturbations smooth

Send offprint requests to: Arhipova N.A.
e-mail: arna@lukash.asc.rssi.ru

out, whereas at large scales ($k < k_{FS}$) they grow to form the gravitationally bounded objects of mass greater than M_{FS} (Novikov & Zel'dovich 1975). Due to the various free-streaming pathes of cold and hot particles ($k_{FS}^{(c)} \gg k_{FS}^{(\nu)}$) the growth rate of density perturbations is different in CDM and HDM components. The linear perturbation power spectrum formed by redshift z, $P(k,z)$, depends strongly on the abundance of each component (Ω_c, Ω_ν) and is given by the production of the initial power spectrum, $P_0(k)$, the transfer function, $T(k,z)$, and the perturbation growth rate, $D(z)$ (e.g. Padmanabhan 1993). Both the transfer function and the growth factor are DM dependent and describe the evolution of initial density perturbations during expansion of the Universe (Zakharov 1979, Bardeen et al. 1986, Holtzman 1989, Eisenstein & Hu 1998, Novosyadly et al. 1999a).

All cosmological models have been re-addressed after CMBA experimental detection (Bennett et al. 1996) to reveal their positive and negative features.

Both ΛCDM and MDM models have met several difficulties.

As far as ΛCDM models are concerned (Kofman et al. 1993, Eke et al. 1996, Liddle et al. 1996a,b, Primack & Klypin 1996) they demand a high value of the cosmological constant ($\Omega_\Lambda \geq 0.7$). In this case ΛCDM is able to fit a set of LSS observational constraints whereas at small scales it overproduces the number of collapsed objects by a factor of 2 in comparison with the corresponding number of gravitationally bounded objects in galactic cataloges (Klypin et al. 1996, Gawiser 2000, Bond et al. 2000).

Regarding MDM models the difficulties are related to late galaxy (quasar) formation (Pogosyan & Starobinsky 1995, Komberg et al. 1996) and too high numbers of obtained galaxy clusters (Ma 1996, Valdarnini et al. 1998, Gardini et al. 1999, Rahman & Shandarin 2001). Standard MDM with one, two or three species of massive neutrino are ruled out at 2σ CL (Novosyadlyj et al. 1999b).

A way to overcome the sMDM difficulties could be the consideration of decaying neutrinos (Bonometto & Pierpaoli 1998), or models with cosmic gravitational waves (CGW) (Arkhipova et al. 1999, Melchiorri et al. 1999), or a non-zero cosmological constant (ΛMDM) (Kahniashvili et al. 1996, Tegmark 1999). In this paper we consider the models with stable neutrinos.

An importance of fundamental CGW has been emphasized by Starobinsky (1979), Rubakov et al. (1982), Lukash & Mikheeva (2000). Theoretical predictions for cluster abundance in MDM models with CGW has been presented by Ma (1996) for the red spectra of density perturbations ($n < 1$), and by Mikheeva et al.(2001) for both blue ($n > 1$) and red ($n < 1$) scalar perturbation spectra. Although ΛCDM models with $\Omega_m \leq 0.3$ normalized by COBE 4-year data (Bunn & White 1997) are consistent with the cluster number density test, in order to archive an agreement with CMBA and cluster abundance data in MDM models it is necessary to take into account the CGW contribution in the derived value of $\Delta T/T$ at 10^0 angular scale. The latter is estimated by parameter T/S, the ratio of the tensor to the scalar mode contributions.

As an alternative to ΛCDM and MDM, the ΛMDM models have been considered by Valdarnini et al.(1998), Novosyadlyj et al.(1999b), Primack & Gross (2000), Andres et al.(2000). The advantage of these models is in retaining the inflationary paradigm and the flat Harrison-Zel'dovich spectrum ($n = 1$) with a smaller value (comparing to ΛCDM) of cosmological Λ-term. Other notable features of ΛMDM are the possibility of negligible CGW contribution (T/S =0) and a small fraction of hot particles: even 10% of massive neutrino in matter content ($\Omega_\nu/\Omega_m \sim 0.1$) could change the value of the cosmological constant, being however in good agreement with other independent tests: CMB data (Hu et al. 2000), distant SNIa (Perlmutter et al. 1999), Hipparcos data (Feast & Catchpole 1997), QSO lensing (Kochanek 1996).

In this paper we consider MDM (with CGW) and ΛMDM (with T/S=0) applying to these models the cluster abundance and evolution tests. The evolution test constrains most efficiently the parameter Ω_m, and hence Ω_Λ in spatially flat cosmological models (e.g. Bahcall et al. 1997, Donahue &Voit 1999, Henry 2000). Our aim is to demonstrate how the presence of hot matter influences the estimation of Ω_Λ, h, and other cosmological parameters. Here, instead of doing an exact χ^2 analysis, we rather look for the tendencies and correlations between cosmological parameters when introducing a fraction of massive neutrino in the Universe.

We describe our models in Section 2, and the galaxy cluster tests in Section 3. The results are discussed in Section 4, with the conclusions in Section 5.

2. Cosmological models with massive neutrino

We assume that DM is given by mixture of CDM and HDM components in the flat background space. The free model parameters are:

- Ω_m, the total matter density in the Universe ($\Omega_m = 1 - \Omega_\Lambda = \Omega_b + \Omega_c + \Omega_\nu$, the latter being density parameters of baryons, cold, and hot particles, respectively);
- $f_\nu \equiv \Omega_\nu/\Omega_m$, the fraction of hot DM;
- N_ν, the number of massive neutrino species;
- h , the Hubble constant in units 100 km s^{-1}Mpc^{-1};
- n , the slope-index of post-inflationary density perturbation power spectrum.

The massive and massless neutrinos[1] are described by the corresponding distribution functions which are evaluated from the Boltzmann-Vlasov collisionless equations.

[1] All neutrinos are active, the total number of neutrino species is three. Accordingly, $N_\nu = 1, 2, 3$.

Cold particles are treated as a pressureless fluid ($p_C = 0$). Baryons and photons are described as an ideal hydrodynamic fluid satisfying the Euler equations of motion. Here we choose the fixed value for baryon density parameter, $\Omega_b = 0.015/h^2$. All DM components interact with each other only gravitationally.

The calculation of the perturbation dynamics (the transfer functions) demands a joint solution of the described self-consistent set of equations. It can be done numerically (Seljak & Zaldarriaga 1996) or analytically in long and short wave regions (Zakharov 1979, Bardeen et al. 1986, Holtzman 1989), the possibility of semi-analitical approximations of some transfer functions has been shown by Eisenstein & Hu (1998), Novosyadlyj et al.(1999a).

Assuming the power low post-inflationary density perturbation spectrum, $P_0(k)\infty k^n$, the final power spectrum of total density perturbations can be expressed as

$$P(k,z) = Ak^n T^2(k,z) \left[\frac{D(z)}{D(0)}\right]^2 , \quad (1)$$

where A is the normalization constant, $T(k,z)$ is the total transfer function, $D(z)$ is the growth factor[2],

$$D(z) = \frac{g(\Omega_m(z))}{1+z} , \quad (2)$$

and $g(\Omega_m(z))$ is the suppression coefficient (Kofman & Starobinsky 1985).

According to Carroll et al.(1992) the function $g(x)$ can be approximated as

$$g(x) = 5x\left(2x^{\frac{4}{7}} + \frac{2+x(209-x)}{70}\right)^{-1} , \quad (3)$$

abd the current matter abundance $\Omega_m(z)$ is written as follows:

$$\Omega_m(z) = \Omega_m \frac{(1+z)^3}{1-\Omega_m+(1+z)^3\Omega_m} , \quad \Omega_m \equiv \Omega_m(0) . \quad (4)$$

We use the transfer function approximations for sCDM (Bardeen et al. 1986) and ΛMDM (Eisenstein & Hu 1998) models.

3. Cluster mass and temperature functions and the evolution test

The mass function for the gravitationally bounded halos of mass greater than M formed in the flat Universe by redshift z is given by (Press & Schechter 1974)

$$N(>M,z) = \int_M^\infty n(M',z)dM' \quad (5)$$

[2] The function $D(z)$ is given as the growth rate of linear density perturbations in the model without hot matter (with the same parameters Ω_m, h, and n).

where $n(M,z)dM$ is the comoving number density of collapsed objects with masses lying in the interval $(M, M+dM)$:

$$n(M,z) = \sqrt{\frac{2}{\pi}}\frac{\rho\delta_c}{M}\frac{1}{\sigma^2(R,z)}\left|\frac{d\sigma(R,z)}{dM}\right| e^{-\frac{\delta_c^2}{2\sigma^2(R,z)}} . \quad (6)$$

$M = \frac{4}{3}\pi\rho R^3$, ρ is the mean matter density, and δ_c is the critical density contrast for a linear overdensity able to collaps. The rms amplitude of density fluctuation in the spheres of radius R at redshift z is related to the power spectrum as

$$\sigma^2(R,z) = \frac{1}{2\pi^2}\int_0^\infty P(k,z)|W(kR)|^2 k^2 dk, \quad (7)$$

where $W(kR)$ is a Fourier component of the top-hat window function: $W(x) = \frac{3}{x^3}(\sin x - x cos x)$. We denote $\sigma_R \equiv \sigma(R,0)$ for $z = 0$.

In the matter-dominated Universe $\delta_c = 1.686$ (Eke et al. 1996, Viana & Liddle 1996), in the flat models with Λ-term δ_c depends weakly on the current matter abundance (Liddle et al. 1996a, Locas & Hoffman 2000).

The theoretical mass functions obtained with the help of the Press-Schechter formalism (eqs.(5,6,7)) are in good agreement with other methods including numerical simulations (Efstathiou et al. 1988, Lacey & Cole 1994, Gross et al. 1998, Eke et al. 1998, Bode et al. 2000, Wu 2000). Due to the exponential dependence of $n(M,0)$ on σ_R (see eq.(6)) the cluster number density is very sensitive to the value of σ_8 .

Rich galaxy clusters are strong X-rays sources characterised by the gas temperature. The physical mechanisms of clusters formation promp the relation between cluster mass and temperature, $T_g\infty M^{2/3}$, confirmed by numerical simulations (e.g. Navarro et al. 1995). The proportionality coefficient depends slightly on cosmological model (mainly, on the parameter Ω_m). Here, we use the $T-M$ relation for isothermal gas given by Eke et al. (1996):

$$T_g = \frac{7.75}{\beta}\left(\frac{6.8}{5X+3}\right) M_{15}^{\frac{2}{3}} \left(\frac{\Omega_m}{\Omega_m(z)}\right)^{\frac{1}{3}} \left(\frac{\Delta_{cr}}{178}\right)^{\frac{1}{3}} (1+z) \text{ Kev} \quad (8)$$

where $\beta(\simeq 1)$ is the ratio of the galaxy kinetic energy to the gas thermal energy, $X(\simeq 0.76)$ is the hydrogen mass fraction, and the mass M_{15} is given in units $10^{15}h^{-1}M_\odot$. The value Δ_{cr} is the ratio of a mean halo density (within the virial radius of collapsed object) to the critical density of the Universe at corresponding redshift. For $\Omega_m \leq 1$, Δ_{cr} can be derived analytically and approximated as $\Delta_{cr} = 178\Omega_m^{0.45}$ for $z = 0$.

As well as the cluster mass function $N(>M) \equiv N(>M,0)$, the X-cluster temperature function $N(>T) \equiv N(>T,0)$ is also sensitive to σ_R. The comparison between the observed temperature (Henry & Arnaud 1991) and mass (Bahcall & Cen 1993) functions shows a good agreement with eq.(8) at $z = 0$. Comparing with observational data both tests provide a powerful constraint on σ-value in different DM models (Valdarnini et al. 1998, Bahcall

& Fan 1998, Durrer & Novosyadlyj 2001). On the other hand CMBA strongly depends on the density perturbation spectrum as well. The consistency with all these tests determines the model parameters (e.g. Bridle et al. 1999, Mikheeva et al. 2001).

The cluster mass (temperature) function and its evolution in ΛCDM has been discussed in detail (e.g. Viana & Liddle 1998, Bode et al. 2000, Henry 2000). To achieve a better consistency with cluster evolution data the value of Ω_m should be lower than that obtained from the cluster number density test at $z = 0$.

The dependence of the cluster mass (temperature) function on z originates due to different growth rate of density perturbations in DM models. It is reflected in the appearance of the suppression factor $g(\Omega_m(z))$ in models with $\Omega_\Lambda \neq 0$ (see eqs.(3,4)):

$$\sigma(R,z) = \frac{\sigma_R}{1+z}\frac{g(\Omega_m(z))}{g(\Omega_m)} \ . \qquad (9)$$

Below, we consider the cosmological models containing massive neutrino. In this case z-dependence is also present in the transfer function due to the free-streaming of hot matter particles (the effective growth rate becomes scale dependent). For the mass estimation of rich galaxy clusters (see eq.(6)), the effect of free-streaming is negligible.



Fig. 1. The present day cluster abundance $N(>T)$ in MDM models normalized by $\sigma_8 = 0.52$ with $\Omega_\nu = 0, 0.1, 0.2, 0.3$ (solid, dot, dash, long-dash lines, resp.), $\Omega_b = 0.015/h^2$, $h = 0.6$, $n = 1$. The needed T/S is shown. The data points by Henry & Arnaud (1991).

4. Results and Discussion

4.1. MDM models with non-zero tensor mode

To test MDM models with a zero cosmological constant we follow the normalization procedure given by Arkhipova et al. (1999), Mikheeva et al. (2001). All models are normalized by σ_8 by the best fit present day cluster mass (Bahcall & Cen 1993, Bahcall et al. 1997) and temperature (Henry & Arnaud 1991, Donahue & Voit 1999) functions. To achieve an agreement with $\Delta T/T$ data (Bennet et al. 1996) the contribution of cosmic gravitational waves is required. The value of the derived parameter T/S depends mainly on the spectral index n, the abundance of hot matter Ω_ν, and the Hubble constant h.

Fig.1 shows the present-day cluster temperature functions $N(>T)$ for different values of Ω_ν in MDM models with CGW normalized by $\sigma_8 \simeq 0.52$. The normalization does not depend on Ω_ν, therefore all curves cross each other at some fixed point corresponding to the mass M in the sphere of radius $8h^{-1}$Mpc.

A significant contribution of both CGW and massive neutrinos is needed to fit the large scale CMBA observational data. The consideration of the red spectra ($n < 1$) cannot reduce T/S substantially. E.g., in MDM models with fixed $\Omega_\nu = 0.2$, $h = 0.6$, and $n = 0.9, 1, 1.1$: T/S=$0.6, 1.7, 3.6$ respectively, which strongly suppresses the height of the first acoustic peak in $\Delta T/T$ spectrum (Mikheeva et al. 2001).

In this respect, higher values of Ω_ν would be preferable for MDM models: an increase of $\Omega_\nu \geq 0.2$ reduces the parameter T/S, however the problem arises with small-scale clustering and the Ly_α-cloud formation tests (Gnedin 1998). On the contrary, at low $\Omega_\nu < 0.2$ neither possible changes of h nor models with three species of massive neutrino can decrease sufficiently a high contribution of CGW into CMBA at large angular scales. The latter obviously leads to a low height of the first acoustic peak inconsistent with the BOOMERANG observations (Netterfield et al. 2001). Fig. 2 demonstrates that the introduction of even a small cosmological constant would drastically improve the situation with MDM-CGW model parameters.

But let us return again to MDM models without Λ-term. Fig. 3 presents the evolution of galaxy clusters. None of the considered models can fit the data at high $z > 0.3$. We varied h (Fig. 3a), Ω_ν (Fig. 3b), and n (not shown). The evolution tracks are practically independent of the post-inflationary spectral index for $n \in (0.9, 1.1)$. Even neglecting the observational problems with Ly_α-forest, the case with high $\Omega_\nu \geq 0.2$ cannot help to achieve an agreement with the evolution of galaxy clusters in MDM dominated models. All the models predict the number of galaxy clusters at high $z \geq 0.3$ at least two orders of magnitude smaller than the observed one. This fact indicates that the evolution of density perturbations should be slower than that found in MDM without a cosmological constant.









Fig. 2. The introduction of a cosmological constant in MDM-CGW models normalized by $\sigma_8 = 0.52\Omega_m^{(-0.52+0.13\Omega_m)}$ with $\Omega_\nu = 0.1$, $\Omega_b = 0.015/h^2$, $h = 0.6$: (a) T/S as a function of n for $\Omega_\Lambda = 0, 0.3, 0.4, 0.5$ (solid, dot, dash, long-dash lines, resp.), and (b) the height of the first acoustic peak, $R^{\frac{1}{2}} = \frac{\delta T_p}{70\mu K}$, as a function of Ω_Λ for $n = 0.9, 0.95, 1, 1.05, 1.1$ (solid, dot, dashed, long-dashed, dashed-dot lines, resp.).

Fig. 3. The cluster evolution $N(> M = 8 \cdot 10^{14} M_\odot, z)$ in MDM models normalized by $\sigma_8 = 0.52$ with $n = 1$, (a) $\Omega_\nu = 0.2$, $h = 0.5, 0.6, 0.7$ (solid, dot, dash lines, resp.), and (b) $h = 0.6$, $\Omega_\nu = 0, 0.1, 0.2, 0.3$ (solid, dot, dash, long-dash lines, resp.). The needed T/S is shown. The data points by Bahcall & Fan (1998).

4.2. MDM models with non-zero cosmological constant

Let us consider flat ΛMDM models without CGW normalized by the COBE 4-year data. This allows an explicit derivation of σ_8(cmb) as a function of model parameters. On the other hand, from observational abundance of galaxy clusters it is possible to obtain σ_8(cl) as a function of Ω_m, e.g. $\sigma_8(\mathrm{cl}) = 0.52\Omega_m^{-0.52+0.13\Omega_m}$ (Eke et al. 1996, Liddle et al. 1996b). The coincidence between σ_8(cmb) and σ_8(cl) would limit the parameter Ω_Λ by ~ 0.7 in ΛCDM (Bahcall & Fan 1998, Henry 2000) and $\sim 0.5 - 0.6$ in ΛMDM (Valdarnini et al 1998, Novosyadlyj et al. 1999b) models.





Fig. 4. The cluster evolution $N(>M=8\cdot10^{14}M_\odot,z)$ in ΛCDM models normalized by $\sigma_8=0.52\Omega_m^{(-0.52+0.13\Omega_m)}$ with $\Omega_\Lambda=0.7$, $\Omega_b=0.015/h^2$, (a) $n=1$, $h=0.65,0.70,0.75$ (dot, dash, long-dash lines, resp.), and (b) $h=0.7$, $n=0.8,0.9,1.0,1.1$ (solid, dot, dash, long-dash lines, resp.). The data points by Bahcall & Fan (1998).





Fig. 5. The present day cluster abundance $N(>T)$ in ΛMDM models normalized by COBE 4-year data with $\Omega_\Lambda=0,0.31,0.45,0.65$ (dot-dash, long-dash, dash, dot lines, resp.), $\Omega_b=0.015/h^2$, $h=0.6$, $n=1$, and $f_\nu=0.1$ (a), 0.2 (b). The data points by Henry & Arnaud (1991).

According to Henry (2000) the cluster evolution test has a stronger upper-limited parameter Ω_m in ΛCDM models than does the cluster abundance test at $z=0$. As an example we demonstrate the evolution of the number density of galaxy clusters of mass $M\geq8\cdot10^{14}M_\odot$ within the comoving radius $1.5h^{-1}$Mpc in ΛCDM model with $\Omega_\Lambda=0.7$, as functions of h (Fig. 4a) and n (Fig. 4b). The cluster evolution is not practically affected by changes of the spectral index. Regarding the parameter h, the models with $h>0.6$ are preferable.

While the requested value of cosmological constant in ΛCDM models is quite high, a small amount of hot particles in ΛMDM models leads to the agreement with observations for a smaller Ω_Λ.

We performed the calculation of functions $N(>T)$ for different parameters Ω_Λ and f_ν (Fig. 5). As we can see,

to achieve an agreement with today cluster temperature function, we need a lower value of cosmological constant in the models with a higher fraction of hot matter. For $f_\nu \in (0, 0.2)$, $h \in (0.6, 0.7)$, $n \in (0.9, 1.1)$, the value of cosmological constant remains in the range (1σ CL):

$$0.35 < \Omega_\Lambda < 0.7 \, . \qquad (10)$$

Taking into account the Ly_α forest test, models with $f_\nu \simeq 0.1$ are preferable. In these case parameter Ω_Λ is limited by $0.45 \leq \Omega_\Lambda < 0.65$ (in agreement with some other results, cf. Valdarnini et al.(1998), Novosyadlyj et al.(1999b), Primack & Gross (2000), Durrer & Novosyadlyj (2001)).

To obtain the redshift evolution functions of galaxy cluster abundance, the normalization of the power spectrum is done according to the present-day cluster concentration, σ_8(cl), which depends on Ω_Λ. At 1σ level the best fit for σ_8(cl) is not sensitive to the hot particle abundance, $h \in (0.6, 0.7)$, and $n \in (0.9, 1.1)$. Figs. 6,7 present the galaxy cluster evolution in ΛMDM models as function of parameters Ω_Λ, f_ν, and h. Changing h and n (within their ranges) does not practically influence the evolution tracks. Increasing f_ν reduces the needed value of Ω_Λ (in agreement with the cluster number density test).

For $h \in (0.6, 0.7)$, $n \in (0.9, 1.1)$, the value of Ω_Λ depends mainly on the fraction of hot matter f_ν. We find the following approximation between the parameters Ω_Λ and $f_\nu \in (0, 0.2)$ from the cluster redshift evolution alone (1σ CL):

$$\Omega_\Lambda + 0.5 f_\nu = 0.65 \pm 0.1 \, . \qquad (11)$$

In connection with this degeneracy two points should be emphasized.

* As we see, the redshift evolution test slightly exaggerates the value of Ω_Λ in comparioson with the present day cluster abundance (cf. eq.(10)). To be in the agreement with both tests the value of the cosmological constant in ΛMDM models should be found as $\Omega_\Lambda \in (0.45, 0.7)$.

* Eq.(11) can also be understood as a proportionality between the values of Ω_m and f_ν: more matter leads to a greater fraction of hot matter and *vice verce*. Such a degeneracy is physically clear and could be straightforwardly observed in LSS tests. E.g., Novosyadlyj' et al. (1999b) indicate the correlation between Ω_m and Ω_ν from the cluster power spectrum and Ly_α data. We guess that a joint LSS analysis will confirm eq.(11) with a better precision.

In conclusion, we stress that the present observational data constrain only the parameter $f_\nu \in (0, 0.2)$, the number of species of massive neutrino ($N_\nu = 1, 2, 3$) remains undetermined.

5. Conclusions

Our main conclusions are as follows:



Fig. 6. The cluster evolution $N(> M = 8 \cdot 10^{14} M_\odot, z)$ in ΛMDM models normalized by $\sigma_8 = 0.52\Omega_m^{(-0.52+0.13\Omega_m)}$ with $\Omega_\Lambda = 0, 0.31, 0.45, 0.55, 0.65$ (solid, dot, dash, dot-dash, long-dash lines, resp.), $\Omega_b = 0.015/h^2$, $h = 0.6$, $n = 1$, and $f_\nu = 0.1$ (a), 0.2 (b). The data points by Bahcall & Fan (1998).

– The introduction of massive neutrino in spatially flat dark matter models affects drastically the estimation of the cosmological parameters from galaxy cluster observational data.

– MDM models with cosmic gravitational waves and a negligible Λ-term do not pass the observational



Fig. 7. The cluster evolution $N(>M=8\cdot10^{14}M_\odot,z)$ in ΛMDM models normalized by $\sigma_8=0.52\Omega_m^{-0.52+0.13\Omega_m}$ with $\Omega_\Lambda=0.55$, $\Omega_b=0.015/h^2$, $n=1$, (a) $h=0.6$, $f_\nu=0,0.1,0.2$ (solid, dot, dash lines, resp), and (b) $f_\nu=0.1$, $h=0.6,0.65,0.7$ (dot, dash, long-dash lines, resp.). The data points by Bahcall & Fan (1998).

constraints: they require a significal contribution of both (i) CGW (T/S $\gtrsim$1) and (ii) hot matter ($f_\nu\gtrsim$0.2). The first condition reduces the first acoustic peak in $\Delta T/T$ spectrum by a level inconsistent with the BOOMERANG measurements, the second condition contradicts the Ly_α-system formation test.

– The introduction of even a small cosmological constant in MDM-CGW models essentially improves the situation with the model parameters, indicating that Λ-term is a more powerful instrument (than T/S) to reconcile theory with observations.

– The cluster abundance and/or evolution tests alone clearly hint at the existence of a non-zero cosmological constant in a set of spatially flat ΛMDM models with a negligible amount of CGW. A rise of T/S cannot cancel the Λ-term. However, the opposite is not true: the estimation of T/S in ΛMDM cosmologies requires better data and a more subtle analysis.

– The presence of a small fraction of massive neutrino, $f_\nu\in(0,0.2)$, (i) reduces the required value of cosmological constant (in comparison with ΛCDM), (ii) is preferable from the point of view of the ly_α forest test, and (iii) does not discriminate the estimate of other cosmological parameters, $h\in(0.6,0.7)$, $n\in(0.9,1,1)$, $N_\nu=1,2,3$.

– For $f_\nu\in(0,0.2)$, $h\in(0.6,0.7)$, $n\in(0.9,1.1)$, the value of cosmological constant remains in 1σ ranges:
  * $\Omega_\Lambda\in(0.35,0.7)$ (from nearby cluster abundance),
  * $\Omega_\Lambda\in(0.45,0.75)$ (from cluster redshift evolution),
  * $\Omega_\Lambda\in(0.45,0.7)$ (from the both tests).

– We find the following approximation between parameters Ω_Λ and $f_\nu\in(0,0.2)$ from the cluster redshift evolution alone:

$$\Omega_\Lambda+0.5f_\nu=0.65\pm0.1\ ,\quad(1\sigma\mathrm{CL}).$$

– If massive neutrinos constitute only $\sim10\%$ of the total DM density ($f_\nu=0.1$) the models with $\Omega_\Lambda\in(0.5,0.65)$ satisfy the observational data best.

It may well be that the DM nature is more complex than just ΛMDM with a negligible amount of CGW. Moreover, even in the framework of this simple model we cannot delimit today the fraction and the number of species of massive neutrino, the accuracy of the available data is still low to do it. Nevertheless, we can assert that the abundance of cosmological massive neutrinos is small, $f_\nu<0.2$, corresponding less than few eV to the neutrino mass. Also, today we can conclude that

* the value of the estimated cosmological constant is very sensitive to even a small fraction of massive neutrinos, and

* the continuing progress in LSS cosmological observations assists in solving the DM problem.

Acknowledgments The work of N.A.A. and V.N.L. was partly supported by RFBR (01-02-16274, 00-15-96639), INTAS (97-1192), and SST Program in Astronomy ("kosmomikrofizika"). The work of T.K. was supported by the COBASE program of USNRC.

References

Andres E., Askebjer P., Bai X., Barouch G., et al., 2000, Nucl.Phys.Proc.Suppl. 91 423 (astro-ph/0009242)
Arhipova N.A., Lukash V.N., Mikheeva E.V., 1999, "Gravitational & Cosmology" 5, 159 (astro-ph/9803328)
Bahcall N.A., Cen R., 1993, ApJ 407, L49
Bahcall N.A., Fan X., Cen R., 1997, ApJ 485, 53
Bahcall N.A., Fan X., 1998, ApJ 504, 1
Bardeen J.M., Bond J.R., Kaiser N., Szalay A.S., 1986, ApJ 304, 15
Bennett C.L. et al., 1996, ApJ 464, L1
Bode P., Bachall N., Ford E.B., Ostriker J.P., 2000, astro-ph/0011376
Bond et al., 2000, astro-ph/0005004
Bonometto S.A., Pierpaoli E., 1998, NewA 3, 391 (astro-ph/9806035)
Bridle S.L. et al., 1999, MNRAS 310, 565
Bunn E.F., White M., 1997, ApJ 480, 6
Carroll S.M., Press W.H., Turner E.L., 1992, A&A 30, 499
Donahue M., Voit G.M., 1999, ApJ 523, L137
Durrer R., Novosyadlyj B.S., 2001, MNRAS 324, 560 (astro-ph/0009057)
Efstathiou G., Frenk C.S., White S.D.M., Davis M., 1988, MNRAS 235, 715
Eisenstein D., Hu W., 1998, ApJ 496, 605
Eke V.R., Cole S., Frenk C.S., 1996, MNRAS 282, 263
Eke V.R., Navarro J., Frenk C.S., 1998, ApJ 503, 569
Fang L.Z., Xiang S.P., Li S.X., 1984, A&A 140, 77
Feast M.W., Catchpole R.W., 1997, MNRAS 286, L1
Gardini A., Bonometto S.A., Murante G., 1999, ApJ 524, 510
Gawiser E., 2000, astro-ph/0005365
Gnedin N.Y., 1998, MNRAS 299, 393
Gross M.A., Samerville J.R., Primack J.R., Borgani S., Girardi M., 1998, Large-Scale Structure: Tracks and Traces, Proc. 12th Potsdam Cosmology Workshop, Potsdam, Sept. 15-19, eds. V. Mueller et al., World Scietific, 171
Henry J.P., 2000, ApJ 534, 565
Henry J.P., Arnaud K.A., 1991 ApJ 372, 410
Holtzman, J.A. 1989, ApJ Suppl. 71, 1
Hu W., Fukugita M., Zaldarriaga M., Tegmark M., 2000, astro-ph/0006436
Jaffe A.H. et al. 2001, Phys. Rev. Lett. 86, 3475
Kahniashvili T.A., B.S. Novosyadlyj, R. Valdarnini, 1996, Phys. Hel. Acta 71, 454
Klypin A., Primack J., Holtzman J., 1996, ApJ 466, 13
Kochanek C.S., 1996, ApJ 466, 638
Kofman L.A., Gnedin N.Y., Bahcall N.A., 1993, ApJ 413, 1
Kofman L.A., Starobinsky A.A., 1985, Sov. Astron. Lett. 9, 643
Komberg B.V., Kravtsov A.V., Lukash V.N., 1996, MNRAS 282, 713
Lacey C., Cole S., 1994, MNRAS 271, 676
Liddle A.R., Lyth D.M., Roberts D., Viana P., 1996a, MNRAS 278, 644
Liddle A.R., Lyth D.M., Viana P., White M., 1996b, MNRAS 282, 281
Locas E.L. , Hoffman Y., 2000, astro-ph/0011295
Lucchin F., Colafrancesco S., de Gasperis G. et al., 1996, ApJ 459, 455
Lukash V.N., 1991, Annals New York Acad. of Sci. 647, 659
Lukash V.N., Mikheeva E.V., 2000, Int. J. of Mod. Phys. A 15, 3783
Ma C.-P., 1996, ApJ 471, 13
Melchiorri A., Sazhin M.V., Shulga V.V., Vittorio N., 1999, ApJ 518, 562
Mikheeva E.V., Lukash V.N., Arkhipova N.A., Malinovsky A.M., 2001, Astron. Zh. 78, 195 (Astron. Reports 45, 163)
Navarro J.F., Frenk C.S., White S.D.M., 1995, MNRAS 275, 720
Netterfield C.B. et al., 2001, astro-ph/0104460
Novikov I.D., Zel'dovich Ya.B., 1975, Structure and Evolution of the Universe, 'Nauka' (in Russian)
Novosyadlyj B.S., Durrer R., Lukash V.N., 1999a, A&A 347, 799
Novosyadlyj B.S., Durrer R., Gottlober S., Lukash V.N., Apunievych S., 1999b, A&A 356, 418
Padmanabhan T. 1993, Structure Formation in the Universe, Cambridge Univ. Press
Peacock J.A., 1997, MNRAS 284, 885
Peebles P.J.E., 1984, ApJ 284, 439
Perlmutter S. et al., 1999, ApJ 517, 565
Pogosyan D.Yu., Starobinsky A.A., 1995, ApJ 447, 465
Press W.H., Schechter P., 1974, ApJ 187, 425
Primack J.R., Klypin A. 1996, Nucl.Phys.Proc.Suppl 51B, 30
Primack J.R., Gross M.A., 2000, astro-ph/0007165
Rahman N., Shandarin S., 2001, astro-ph/0010228
Rubakov V.A., Sazhin M.V., Verjaskin A.M., 1982, Phys. Lett. B 115, 189
Seljak U., Zaldarriaga M., 1996, ApJ 469, 437
Shafi Q., Stecker F.W. 1984, Phys. Rev. Lett. 53, 1292
Smoot G.F., Bennett C.L., Kogut A. et al. 1992, ApJ 396, L1
Starobinsky A.A., 1979, JETF Lett 30, 719
Tegmark M., 1999, ApJ 514, L69
Valdarnini R., Bonometto S.A., 1984, Phys. Lett. A 103, 369
Valdarnini R., Kahniashvili T.A., Novosyadlyj B.S., 1998, A&A 311, 1
Viana P.T.P., Liddle A.R., 1996, MNRAS 281, 323
Viana P.T.P., Liddle A.R., 1998, MNRAS 303, 535
Wu J.-H.P., 2000, astro-ph/0012207
Zakharov A.V., 1979, JETP 77, 434 (in Russian)